
Journées Scientifiques Annuelles d'Arbo-France

Moyens innovants de lutte contre les vecteurs d'arbovirus

23-24 juin 2025, Val de Grâce - Paris

Overview du contrôle de la transmission des arboviroses liées à Aedes

Historique et bilan du contrôle des Aedes

par Fabrice Chandre

Historique et bilan de la lutte contre les Aedes

23-24 Juin 2025

Fabrice CHANDRE



Maladies Infectieuses et Vecteurs
Écologie, Génétique, Évolution et Contrôle



**RÉPUBLIQUE
FRANÇAISE**


*Liberté
Égalité
Fraternité*



Institut de Recherche
pour le Développement
FRANCE



Succès historiques du contrôle d'*Aedes aegypti*

- ▶ Cuba (1901-1912), Canal de Panama (1904-1913), W. Gorgas: réduction de fièvre jaune par contrôle larvaire (élimination gîtes, huiles minérales) + isolement des malades, contraventions
- ▶ Brésil (1930-1942) F. Soper: réduction fièvre jaune et élimination partielle d'*Aedes aegypti*
- ▶ Am. Centrale, Am. Sud, Caraïbes (1946-1970) contrôle larvaire & traitement périfocal DDT- élimination d' d'*Aedes aegypti* dans 23 pays



Recolonisation par *Aedes aegypti*

- ▶ Réduction des ressources pour le contrôle des vecteurs (*Aedes*, *Anopheles*) vers autres priorités de santé publique à partir de 1970
- ▶ Accroissement de la population, urbanisation non contrôlée, transport marchandises et personnes
- ▶ Résistance au DDT



The emergence of dengue hemorrhagic fever in the Americas associated with introduction of new virus serotypes and the development of hyperendemicity.


Gestion Intégrée des Vecteurs (OMS, 2004)

- ▶ processus rationnel de prise de décisions dans la perspective d'une utilisation optimale des ressources affectées au contrôle des vecteurs

Améliorer l'efficacité et l'efficience des interventions de contrôle des vecteurs

Réduire l'impact écologique et le recours aux insecticides

Réduire le nombre d'interventions en agissant sur plusieurs vecteurs et/ou plusieurs maladies vectorielles



Aedes aegypti et *Aedes albopictus*

- ▶ Très grande diversité des habitats larvaires

- ▶ Gîtes domestiques



- ▶ Gîtes périurbains



- ▶ Gîtes naturels



- ▶ Plasticité physiologique et écologique des *Aedes*

- ▶ Anthropophile mais opportuniste (anx domestiques)

- ▶ Diurne, pique au lever du jour et en fin d'après midi

- ▶ La femelle pond qq œufs dans de nombreux gîtes


- ▶ En l'absence de pluies les œufs résistent à la dessiccation

- ▶ Diapause des œufs en hiver pour *A. albopictus*



Lutte intégrée contre les *Aedes*

- ▶ Surveillance entomologique (indices larvaires, piégeage adultes)
- ▶ Protection réserves d'eau
élimination petits gîtes
- ▶ Information, éducation
sanitaire, mobilisation sociale



Lutte intégrée contre les *Aedes*

► Lutte larvicide: chimique, biologique




Mesocyclops ruttneri
M. aspericornis
M. thermocyclopoides
M. affinis
M. woutersi
M. ogunnus

► Lutte adulticide: thermonébulisation ou ULV en cas de circulation virale



► Protection personnelle: vêtements longs, imprégnés, répulsifs, moustiquaires




Insecticides en lutte antivectorielle

► Principales familles d'insecticides utilisées en LAV

- Pyréthrinoïdes (et DDT Organochloré)
 - Inhibiteurs du Canal Sodium Voltage dépendant
 - Action rapide (knockdown), effet irritant, adulticides
- Organophosphorés et Carbamates
 - Inhibiteurs de l'acétylcholinestérase, adult-/larvicides
- Régulateurs de croissance
 - Analogues hormonaux, perturbation du développement, larvicides
- Insecticides d'origine bactérienne (*Bti*, *Bs*)
 - Toxique/ingestion, épithélium intestinal, larvicides


94% agissent
sur 2 cibles !




% tonnage annuel 2010-2019 pour
le contrôle des vecteurs d'arbovirus

WHO, 2021 Global Insecticide Use for
Vector-Borne Disease Control, 6th Ed.

Principaux mécanismes de résistance





Résistance d'*Aedes aegypti* à la deltaméthrine (adulticide) (*Moyes et al., 2017, Plos NTD*)




Fig 2. The frequency of resistance to deltamethrin in *Ae. Aegypti*, 2006–2015. Adult bioassays using 0.05% insecticide for 1 hour are denoted as circles and results from nonstandard adult bioassays (including different diagnostic doses and exposure periods) are denoted as triangles. The map is zoomed to the 3 regions with data. (A) Americas. (B) Africa/Arabian Peninsula. (C) Asia.

Résistance d'*Aedes aegypti* au téméphos (larvicide) (*Moyes et al., 2017, Plos NTD*)






Fig 3. The level of *Ae. aegypti* resistance to temephos, 2006–2015. The ratio of the lethal concentration required to kill half of the sample (LC₅₀ value) obtained by each study to the value obtained for the Rockefeller susceptible strain across studies was calculated. The ratios were then split into 5 classes: values less than 2-fold higher than Rockefeller and each quartile of the remaining distribution. The map is zoomed to the 3 regions with data. (A) Americas. (B) Africa. (C) Asia.

Canal sodium et mutations *kdr* chez *Aedes aegypti*



Moyes et al. 2017
Plos NTD



V1016G, F1534C aussi présentes chez *Ae. albopictus*

+ V410L
Haddi et al. 2017

Fig 5. The geographical distribution of the 10 known voltage-gated sodium channel (VGSC) mutations in *Aedes aegypti* across the 3 continents in which they have been detected. Association of each mutation with pyrethroid resistance is shown in the key. Font size gives an indication of relative frequency.

Enzymes impliquées dans la résistance métabolique (validation fonctionnelle) – Anses 2021

Tableau 10. Principaux marqueurs moléculaires utilisables pour détecter la résistance métabolique


Espèce	Gène ¹	Famille Enzymatique	Insecticides	CNV détecté ²	Références
<i>Ae. aegypti</i>	CYP9J32	P450	PYR	oui	Stevenson <i>et al.</i> 2012, Kasai <i>et al.</i> 2014, Faucon <i>et al.</i> 2015, Smith <i>et al.</i> 2016*, Moyes <i>et al.</i> 2017*, Faucon <i>et al.</i> 2017, Cattell <i>et al.</i> 2020a, Cattell <i>et al.</i> 2020b
	CYP9J26	P450	PYR	oui	
	CYP9J28	P450	PYR	oui	
	CYP9J24	P450	PYR	oui	
	CYP9M6	P450	PYR	oui	
	CYP6BB2	P450	PYR	oui	
	CCEAE3A	COE	OP	oui	
	GSTE2	GST	PYR, DDT	oui	
<i>Ae. albopictus</i>	CYP6P12	P450	PYR	non	Grigoraki <i>et al.</i> 2015
	CCEAE3A	COE	OP	oui	Ishak <i>et al.</i> 2016
<i>Cx. pipiens / quinquefasciatus</i>	CYP9M10	P450	PYR	oui	Weill <i>et al.</i> 2000 Hardstone <i>et al.</i> 2010, Scott <i>et al.</i> 2015*, Gong <i>et al.</i> 2017,
	CYP6AA7	P450	PYR	non	
	CYP4H24	P450	PYR	non	
	CPIJ013917 (Est2)	COE	OP	oui	
	CPIJ013918 (Est3)	COE	OP	oui	

¹ Seules les enzymes dont la capacité à dégrader les insecticides a été validée sont indiquées.

² Variations du nombre de copies (CNV) détectées dans les populations naturelles (résistance éventuellement détectable sur ADN génomique). * articles de synthèse.




Impact des pulvérisations spatiales sur *Ae. aegypti* résistant à la deltaméthrine en Martinique (phase II)




Impact de la résistance sur l'efficacité opérationnelles des traitements en Martinique


• Treated arms



• 3 rounds of space spraying, 1 every 3 days



Femelles capturées par pièges BG ext. avant et après traitement



Mortalités relevées en cages sentinelles

Perte d'efficacité des traitements

- En conditions de terrain :
 - Perte totale d'efficacité des traitements ULV à la deltaméthrine sur *Aedes aegypti* à la Martinique

Impact épidémiologique de la résistance

Pas de preuve directe d'un impact épidémiologique de la résistance mais une relation résistance et incidence des cas est suspectée



Incidence microcéphalies attribuables au virus Zika (2016)

Cas de microcéphalies

Résistance temephos (OP)



Incidence microcéphalies plus élevée dans les zones où *Ae. aegypti* est résistant au téméphos


Chediak et al., 2016



Incidence Chikungunya (2007 & 2017)

>200 cas autochtones en **Emilie Romagne** en 2007

~500 cas autochtones cases dans le **Latium** en 2017



Incidence chikungunya plus élevée dans les zones où *Ae. albopictus* est résistant aux pyréthrinoides

Pichler et al., 2019

Bilan et Conclusion

- ▶ M. Chan, Directrice Générale de l'OMS en 2016 (66th WHA)

“Above all, the spread of Zika, the resurgence of dengue, and the emerging threat from chikungunya are the price being paid for a massive policy failure that dropped the ball on mosquito control in the 1970s”

- ▶ Les campagnes préventives de contrôle des Aedes ont été peu à peu remplacées par des réponses d'urgence face aux épidémies
- ▶ Incertitudes sur l'efficacité des méthodes de lutte en fonction du contexte épidémiologique du fait de l'absence d'évaluation robuste
- ▶ Résistance aux insecticides



Bilan et Conclusion

- ▶ Nécessité de développer des méthodes alternatives, insecticides à nouveau mode d'action, outils non insecticides
- ▶ Vector Control Advisory Group (OMS) nouveaux outils doivent démontrer leur impact en terme de santé publique i.e. impact épidémiologique et entomologique via ERC
 - ▶ Répulsifs spatiaux (Pérou, Sri Lanka)
 - ▶ Piège Autodissémination (Laos, Cambodge, Malaisie)
 - ▶ Wolbachia IP (Indonésie, Brésil, Vietnam)
 - ▶ Wolbachia TII + TIS (Singapour)
 - ▶ TIS (Polynésie, non ERC)



Overview du contrôle de la transmission des arboviroses liées à Aedes

Evaluation et Impacts des méthodes de contrôle

par David Roiz



Evaluation and impacts of vector control methods

David Roiz



MIVEGEC Unit. Institut de Recherche pour le Développement (IRD)

23-24 June 2025, ARBOFRANCE, Paris




Menu for today

1. *Aedes*-borne diseases: limitations on vector control
2. The Integrated *Aedes* Management (IAM) framework
3. Integrated Surveillance and integrated vector control
4. Evidence-based, what evidences we have for several control methods?
5. Pitfalls for evaluation and creating evidence-based decisions
6. Social mobilisation, intersectoral collaboration, additional activities
7. The rising global economic costs of *Aedes*-borne diseases
8. Multi-sectoral impacts
9. Prospects for the future and take-home messages

The spread of *Aedes* mosquitoes..


Aedes aegypti



Vector control is the mainstay of the prevention
(in research for efficient vaccines)



$$V_c = \frac{ma^2 b p^n}{-\ln(p)}$$

Aedes albopictus




..cause a global health burden: *Aedes*-borne diseases


Dengue




Zika



Yellow fever




Chikungunya




Also in Europe...


Local-acquired **dengue** cases



Imported dengue cases



Chikungunya



Past and present virus
 transmission
 ■ CHIKV
 ■ DENV
 ■ ZIKV
 ■ CHIKV & DENV
 ■ CHIKV & ZIKV
 ■ DENV & ZIKV
 ■ CHIKV, DENV & ZIKV

Heterogeneity of epidemiological
and entomological contexts!

Aedes aegypti

Aedes albopictus




■ Yellow fever mosquito
 ■ Asian tiger mosquito
 ■ Asian tiger and yellow fever mosquito
 ■ Polynesian tiger mosquito

Limitations of *Aedes* vector control

Ineffective implementation (difficulties for top-down)
Inadequate human, financial and infrastructural capacities in low-income
Lack of community engagement
Aversion to insecticiding and insecticide resistance
Lack of political will

Weak evidence base for the effectiveness, costs,
and public health value of *Aedes* vector control strategies




Background – Basis of the IAM framework

REVIEW

Integrated *Aedes* management for the control of *Aedes*-borne diseases

David Roiz^{1*}, Anne L. Wilson², Thomas W. Scott³, Dina M. Fonseca⁴, Frédéric Jourdain¹, Pie Müller^{5,6}, Raman Velayudhan⁷, Vincent Corbel^{1*}

Locally-adapted practical guidance on how and when to deploy vector management in different settings




Scenarios for the Integrated Aedes Management (IAM) framework



These risk scenarios evolved with a non-linear transition!


Sporadic transmission - irregular, locally and unpredictably

Locally established – Aedes in less than 25 km²




Ae. albopictus in Europe

S₄ *Endemic transmission – limited in space but not in time – dengue transmission season*



Bangkok (1980-2010)


S₅ *Epidemic transmission – limited in space and time – excess of normally expected cases*



Zika (2016)




Entomological surveillance



Evaluation of interventions is critical!


Integrated vector control



- In routine or emergency
- Proactive rather than reactive
- Both larvae and adults vs prevalent paradigm targeting larvae
- Non-insecticide based tools

But...

What is the strength of the evidence?




What is the strength of the evidence?

Evidence-based vector control? Improving the quality of vector control trials

Anne L. Wilson¹, Marleen Boelaert², Immo Kleinschmidt³, Margaret Pinder^{1,4}, Thomas W. Scott^{5,6}, Lucy S. Tusting⁷, and Steve W. Lindsay¹

HOW TO DESIGN VECTOR CONTROL EFFICACY TRIALS

Guidance on phase III vector control field trial design
provided by the Vector Control Advisory Group



- **Randomised controlled** trial (individual or cluster randomised)
- **Randomised** cross-over study, step-wedge design, controlled before-and-after study, controlled time series or controlled interrupted time series
- **Non-randomised** cross-over study, step-wedge design, controlled before-and-after study or controlled interrupted time series
- **Case-control, cohort** or **cross-sectional** study
- **Non-randomised controlled trial** or **non-randomised controlled time series**
- Studies **without a control group** or using a **historical control group**

What is the strenght of the evidence?

Evidences

Examples of Epidemiological evidences of Integrated Aedes management

EVIDENCE-BASED DECISION MAKING

- Epidemiological CRCT evidences for [Community source reduction IVM](#) Nicaragua and Mexico (Andersson et al., 2017)
- Epidemiological CRCT evidences for [Targeted Indoor Residual Spraying](#) Mexico and Australia (Manrique-Saide et al., 2020)
- Epidemiological CRCT evidences for [Wolbachia population replacement](#) Indonesia (Utarini et al., 2021)
- Epidemiological CRCT evidences for [Spatial repellent in Peru](#) (Morrison et al., 2022)
- Epidemiological observational evidences for [Space Spraying in Peru](#) (Stoddard et al., 2014)
- Epidemiological observational evidences for [Mass-trapping](#) in Puerto Rico (Barrera et al., 2019)
- Epidemiological observational evidences for [Insecticide treated materials](#) (Bowman et al., 2016)
- Epidemiological observational evidences for [Larviciding Pyriproxifen](#) (Maoz et al., 2017)



What is the strenght of the evidence of innovative methods?

Wolbachia population replacement

The NEW ENGLAND
JOURNAL of MEDICINE


ESTABLISHED IN 1812

JUNE 10, 2021


VOL. 384 NO. 23

Efficacy of Wolbachia-Infected Mosquito Deployments for the Control of Dengue

A. Utarini, C. Indriani, R.A. Ahmad, W. Tantowijoyo, E. Arguni, M.R. Ansari, E. Supriyati, D.S. Wardana, Y. Meitika, I. Ernesia, I. Nurhayati, E. Prabowo, B. Andari, B.R. Green, L. Hodgson, Z. Cutcher, E. Rancès, P.A. Ryan, S.L. O'Neill, S.M. Dufault, S.K. Tanamas, N.P. Jewell, K.L. Anders, and C.P. Simmons, for the AWED Study Group*





Sterile Insect technique (SIT) and Incompatible insect technique (IIT) Meta-analysis: SIT- *Ae. albopictus*: egg reduction of 41.5% (19.4-51.6)



Mass-trapping


Current evidences of the efficacy of mosquito mass-trapping interventions to reduce *Aedes aegypti* and *Aedes albopictus* populations and *Aedes*-borne virus transmission

Ali Jaffal¹, Johanna Fite^{1*}, Thierry Baldet^{2*}, Pascal Delaunay^{3*}, Frédéric Jourdain^{4,5*}, Ronald Mora-Castillo^{6*}, Marie-Marie Olive^{2,5*}, David Roiz⁵



Variability of *Ae. albopictus* female reduction of 25-64 %

Integrated vector control



Aedes albopictus – very limited entomological evidences

ORS: Outdoor Residual Spraying (vs ULV)

Source reduction and door-to-door

Larviciding (Cold-fogging Bti)

Mass-trapping (Gravid traps and adults)

SIT/Wolbachia

Auto-dissemination...

Effectiveness of a multiple intervention strategy for the control of the tiger mosquito (*Aedes albopictus*) in Spain

Gisela Chebab Abramides^a, David Roiz^b, Raimon Guitart^c, Salvador Quintana^d, Irene Guerrero^e, Nuria Giménez^{c,f,*}

1. What is exactly the research question?
2. Epidemiological outcomes (Phase III): dengue cases, serologies, seroincidence
3. Entomological outcomes: BG-Traps, HLR, ovitraps, hatching rate. Never larval indexes
4. Contemporary controls: is always possible? What is a comparison area?
5. Pre- and post-treatment measures
6. Size of the areas / buffer area?
7. Adult contamination effects / trap locations
8. Replications: number of clusters?
9. Power analysis
10. Randomization
11. Statistical analysis
12. Cost and cost-effectiveness
13. Publish positive and Negative results





Figure 1. Schematic of two generic clusters demonstrating the 'fried-egg design'. The diagonal line patterns represent the 'buffer zones'.



*If it's fast and cheap it won't be good.
If it is cheap and good, it will not be fast.
If it is fast and good, it will not be cheap!*



*In endemic areas, we can be cost-effective if we reduce *Aedes* populations by 70-90%, or even 50-70%**

1. Strength of the evidence of the effectiveness (entomological, epidemiological)
2. Economic costs and cost-effectiveness
3. Temporality/sustainability
4. Quality of delivery (coverage, delivery measures)
5. Environmental impact (non-target fauna)
6. Acceptability/legislation
7. Scenarios/ emergence vs routine / *aegypti* vs *albopictus* / DEN vs ZIKA/CHIK

Social mobilisation



- Human behaviour is a common denominator of control programmes
- Epidemiological and entomological evidences for community based control programs
- Communication for behavioural impact (COMBI)
- Adapted to scenario, context, socio-economic, culture and proactive



Multisectoral collaboration and supporting activities



- Inter-sectoral and inter-agency collaborations
- Ministries, NGOs, UN agencies, private sector, gender, religious and academic institutions
- As provision of reliable piped water, solid waste management.

A schematic guide for multisectoral collaboration

Ministry or organisation	Scenario	Activity	Rationale
Ministry of Public works and municipal authorities	S1, S2, S3 and S4	Provision of reliable piped water to households. Provision of sewerage connections. Solid waste management and disposal. Design and maintenance of street storm water drainage systems that do not harbour immature vectors.	Removal of potential <i>Aedes</i> mosquito larval development habitats can reduce adult numbers and virus transmission rates.
Ministry of Housing	S3 and S4	Develop and enforce housing and building codes mandating installation of screening, dependable water supply, waste management and disposal and rainwater runoff control in new housing developments.	Reduce biting of humans by installing screens to prevent entry to houses and buildings. Reduce standing water to prevent immature development.
Ministry of Education	S3 and S4	Incorporate information on <i>Aedes</i> -borne diseases, vectors, transmission and prevention into school curricula and teach hygienic behaviour.	Empower children with knowledge and skills to reduce mosquito populations and virus transmission.
	S3, S4 and S5	Participation of school children in larval surveys, source reduction and larviciding.	
Ministry of Tourism	S3 and S4	Reduce aquatic habitats in and around hotels, community gathering places, markets, etc.	Reduce virus transmission.
	S5	Communicate rapidly to holiday	Supports rapid implementation of

SUPPORTING ACTIVITIES

Capacity building



Advocacy



Research



Policies & laws




MULTISECTORAL
COLLABORATION



Integrated *Aedes* management for the control of *Aedes*-borne diseases

David Roiz^{1*}, Anne L. Wilson², Thomas W. Scott³, Dina M. Fonseca⁴,
Frédéric Jourdain¹, Pie Müller^{5,6}, Raman Velayudhan⁷, Vincent Corbel^{1*}



Aedes-borne diseases they also cause a global economic burden

The economic burden needs to be appropriately quantified to facilitate the management decisions. But the **economic impacts have not been sufficiently studied**

Since 2016, the InvaCost initiative has generated a database of the reported costs of biological invasions, including mosquitoes. *Ae. aegypti* and *Ae. albopictus* among **the costliest invasive species worldwide**

High and rising economic costs of biological invasions worldwide

nature

<https://doi.org/10.1038/s41586-021-03405-6>

Received: 8 April 2020

Christophe Diagne¹, Boris Leroy², Anne-Charlotte Vaissière¹, Rodolphe E. Gozlan², David Roiz⁴, Ivan Jarić^{5,6}, Jean-Michel Salles⁷, Corey J. A. Bradshaw⁸ & Franck Courchamp¹✉



SCIENTIFIC DATA



ARTICLE

Received 13 Feb 2016 | Accepted 18 Aug 2016 | Published 4 Oct 2016

Massive yet grossly underestimated global costs of invasive insects


Corey J.A. Bradshaw^{1,2}, Boris Leroy^{1,3}, Céline Bellard^{4,5}, David Roiz^{5,6}, Céline Albert^{1,4}, Alice Fournier¹, Morgane Barbet-Massin¹, Jean-Michel Salles⁵, Frédéric Smard⁵ & Franck Courchamp^{1,7,8}

OPEN

InvaCost, a public database of the economic costs of biological invasions worldwide

C. Diagne¹, B. Leroy², R. E. Gozlan², A.-C. Vaissière¹, C. Assailly^{1,4}, L. Nuninger^{1,4}, D. Roiz⁴, F. Jourdain^{5,6}, I. Jarić^{5,6} & F. Courchamp^{1,7,8}

So, we focused on a more complete research of **REPORTED** costs for both *Aedes* mosquitoes and *Aedes*-borne diseases





The rising global economic costs of *Aedes*-borne diseases and *Aedes* mosquitoes



Contents lists available at [ScienceDirect](#)

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv



The rising global economic costs of invasive *Aedes* mosquitoes and *Aedes*-borne diseases

David Roiz^{a,b,*}, Paulina A. Pontifes^{a,b,1}, Frédéric Jourdain^{a,c}, Christophe Diagne^e, Boris Leroy^f, Anne-Charlotte Vaissière^{d,g}, María José Tolsá-García^{a,b}, Jean-Michel Salles^h, Frédéric Simard^a, Franck Courchamp^d

^a MIVEGEC, Univ. Montpellier, IRD, CNRS, Montpellier, France

^b International Joint Laboratory ELDORADO, IRD/UNAM, Mexico

^c Santé Publique France (French National Public Health Agency), Montpellier, France

^d CNRS, AgroParisTech, Écologie Systématique et Évolution, Université Paris-Saclay, Gif-sur-Yvette, 91190, France

^e CBGP, Université de Montpellier, IRD, CIRAD, INRAE, Institut Agro, 34988 Montpellier-sur-Les, France

^f Unité Biologie des Organismes et Écosystèmes Aquatiques (BOREA, UMR 7209), Muséum national d'Histoire naturelle, Sorbonne Université, Université de Caen Normandie, CNRS, IRD, Université des Antilles, Paris, France


^g ECOBIO (écosystèmes, biodiversité, évolution) - UMR 6553, CNRS, Université de Rennes, 263 Avenue du Général Leclerc, 35042 Rennes, France

^h CEE-M, Univ. Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France

HIGHLIGHTS

- Dengue, Zika and chikungunya are transmitted by *Aedes aegypti* and *Aedes albopictus*.
- These invasive species carry a significant but not well-characterized economic cost.
- Our study reports costs from 166 countries and territories, spanning 45 years.
- The cumulative reported cost amounted to at least US\$ 94.7 billion.
- Costs are increasing and only a modest proportion (1/10) is invested in prevention.

GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: Rafael Mateo

Keywords:

Economic costs
Aedes aegypti
Aedes albopictus
Dengue
Zika

ABSTRACT

Invasive *Aedes aegypti* and *Aedes albopictus* mosquitoes transmit viruses such as dengue, chikungunya and Zika, posing a huge public health burden as well as having a less well understood economic impact. We present a comprehensive, global-scale synthesis of studies reporting these economic costs, spanning 166 countries and territories over 45 years. The minimum cumulative reported cost estimate expressed in 2022 US\$ was 94.7 billion, although this figure reflects considerable underreporting and underestimation. The analysis suggests a 14-fold increase in costs, with an average annual expenditure of US\$ 3.1 billion, and a maximum of US\$ 20.3 billion in 2013. Damage and losses were an order of magnitude higher than investment in management, with

A pionnering and exhaustive global study of reported costs
166 countries and territories covering **45 years**
Data to **inform evidence-based policy** into the nature
and distribution of *Aedes*-borne costs.

PRISMA Methodology

Methodology to **avoid double-counting**

Reported evidence from high-reliability records



Data
Compilaton



Data
Extraction



Data
treatment



Data
analysis



DRYAD



Who we :

The rising global economic costs of invasive *Aedes* mosquitoes and *Aedes*-borne diseases



 What are the <u>types of costs</u> and <u>who paid</u> for them?				
	Damage	Management	Losses	Non-market costs
Health Providers (public/private)	Direct medical costs Non-medical costs	Surveillance Vector control Prevention Blood and organ screening Vaccination		
Individuals	Out-of-pocket costs Indirect costs (wage losses, debt)	Personal protection		Quality of life
Community	Indirect costs lost productivity, disability benefits	Research and funding Community-based vector control	Tourism Trade losses (commercial) Economic growth	

Massively increasing reported global costs

- The **minimum** estimate of reported cumulative costs is \$94.7 billion (1975-2020)
- The average **annual** cost is \$3.1 billion, with a peak of \$20.3 billion in 2013
- Costs **multiplied by 14** in 2 dengue decades (1995 to 2014)
- The costs of the longer-term consequences of chikungunya and zika (microcephaly, osteoarthritis, etc.) are estimated at **\$318 billion**.
- Costs are largely **underestimated and underreported**




Geographical breakdown of reported costs



Within the same order of magnitude as other large-scale global estimates (Shepard et al., 2016; Selck et al., 2014; World Bank, 2018)

Geographical breakdown of reported costs

Chikungunya




Costs in US\$ millions



Geographical breakdown of reported costs


Both *Aedes* species





Costs in US\$ millions



How do the costs increase in relation to cases?







Who pays?




- Healthcare providers absorb the majority of direct medical costs due to dengue
- Indirect medical costs (after-effects, productivity, etc.), which occur over the long term and are more difficult to quantify, are borne by individuals and the community
- The losses borne mainly by the community (tourism, trade, economic growth) are difficult to quantify
- A disproportionate burden among impoverished population without access to public health services, reinforcing the poverty trap

Relationship damage vs management

- Expenditure on damage and losses is 10 times higher than the investment in management, surveillance and control.
- Only 10% has been allocated to management and prevention



Relationship damage vs management



1 square = 1 million US\$



Damage
86%




Management
7.6%

A limited investment in mosquito control
Potential benefits are the avoided costs
Effective control actions could reduce damage
in a long-term


Early action with proactive (preventive) strategies rather
than reactive (emergency) responses or even inaction

Effective, sustainable and integrated management would
be beneficial for public health and human societies


The magnitude of the relationship depends –in part- on the scenario



We can have a preliminar idea based on the differences in magnitudes, but the benefits of avoiding cases are at a short- and at a long- term



Example of France 1990-2020



Multi-sectoral impacts



Moustique tigre en France hexagonale : risque et impacts d'une arbovirose

Avis de l'Anses
Rapport d'expertise collective

Juillet 2024

GTARBO

Marie-Claire Paty , Bénédicte Apouey , Clémentine Calba , Éric Cardinale , Carine Milcent , David Roiz , Elodie Rouviere , Jean-Michel Salles , Frédéric Simard, Yannick Simonin, Josselin Thuilliez, Émeline Barrès , Véronique Raimond , Johanna Fite, Florence Étoré, Karine Fiore

Take home messages

- To strengthen robust evidences for innovative non-insecticide approaches
 - Entomological and epidemiological outputs and risk thresholds
- Cost-effectiveness / temporality / acceptability/ quality of delivery
 - Sustainability/Proactive/Synergies/ No magic bullets
 - Locally-adapted/ focus on high risk areas (hot-spots)
 - Highly increasing underreported economic costs
 - Limited investment in vector control campaigns
- Multi-sectoral impacts on the society, media, politics, losses...

We need more operational research to foster evidence-based vector control guidance tailored to local settings and available resources



david.roiz@ird.fr



Institut de Recherche pour le Développement (IRD)-MIVEGEC.

Verified email at ird.fr - [Homepage](#)


Ecology of mosquito-borne ... Aedes-borne diseases control Aedes albopictus
emerging diseases



Cited by

[VIEW ALL](#)

	All	Since 2020
Citations	7702	5692
h-index	42	36
i10-index	65	63



Public access

[VIEW ALL](#)

1 article

49 articles



Overview du contrôle de la transmission des arboviroses liées à Aedes

Aedes albopictus management: some Italian experience

par Romeo Bellini



**Centro
Agricoltura
Ambiente**
Giorgio Nicoli

Aedes albopictus management: some Italian experience

Romeo Bellini

Centro Agricoltura Ambiente «G. Nicoli»



**Journées scientifiques :
Moyens innovants de lutte contre les vecteurs d'arbovirus.**

Institut Pasteur, Paris

23 & 24 juin 2025



MINISTRY OF HEALTH

National plan for prevention, surveillance and answer to arboviruses 2020-2025

WEST NILE

USUTU

CHIKUNGUNYA

DENGUE

ZIKA

TBE

TOSV




Regione del Veneto

Il portale della Regione del Veneto



REGIONE AUTONOMA
FRIULI VENEZIA GIULIA



REGIONE
PIEMONTE

Emilia-Romagna Regional Plan on arboviruses



ZANZARATIGREONLINE

CHI FA COSA

MONITORAGGIO

NEWS e EVENTI

ZANZARE e VIRUS

COMUNICAZIONE

APPROFONDIMENTI

ZANZARE E ALTRI INSETTI impara a difenderti

Zanzare tigre, zanzare comuni e pappataci possono trasmettere malattie anche gravi.


Proteggi te stesso e i tuoi familiari: contrasta la diffusione di questi insetti, con alcune semplici azioni.



ALBOPICTUS MANAGEMENT PLAN EMILIA-ROMAGNA

- **1. Public health risk assessment**
- **2. Monitoring by ovitraps**
- **3. Standard control measures in public areas**
- **4. Quality control on larvicide treatments in public road drains**
- **5. Emergence control measures in case of DEN, CHIK & ZIKA imported cases detection**
- **6. Community participation**
- **7. Door-to-door control measures in private areas**
- **8. Resistance prevention**

Imported & autochthonous Dengue, Chikungunya, Zika cases in Emilia-Romagna




IMMEDIATE REACTION IN CASE OF IMPORTED CASES DETECTION CHIK / DENGUE / ZIKA

- immediate adulticide treatment in public areas (for three consecutive nights)
- immediate door-to-door larvicide, adulticide and source removal
- area of 100-200 m ray




Autochthonous dengue outbreak in Marche Region, Central Italy, August to October 2024

A. Epidemic curve (n = 199)



Premises Demographic Survey (PDS)



Container typology	No. containers /premise	No. positive containers /premise	No. pupae/ha
Private Catch basin	2.35 ± 2.96	0.33 ± 0.96	237.7 ± 152.46
Plant saucer	5.79 ± 11.75	0.08 ± 0.57	1.72 ± 1.67
Drums	0.09 ± 0.62	0.03 ± 0.27	6.09 ± 11.13
Buckets	0.31 ± 1.16	0.03 ± 0.27	2.08 ± 2.35
Tarpaulins	0.09 ± 1.06	0.00 ± 0.03	0.01 ± 0.05
Tires	0.23 ± 9.75	0.00 ± 0.11	0.13 ± 0.44
Bathtubs	0.02 ± 0.23	0.00 ± 0.05	0.36 ± 0.87
Containers < 1 L	0.28 ± 3.01	0.02 ± 0.20	0.54 ± 0.64
Containers 1-10 L	0.29 ± 1.07	0.03 ± 0.27	2.28 ± 4.80
Containers > 10 L	0.09 ± 0.66	0.01 ± 0.14	1.25 ± 1.89
Public catch basins			196.4 ± 446.73

STANDARD CONTROL MEASURES IN PUBLIC AREAS

- **Regular larval control in public drains
(1.3 MIO)**



- **Mayor's ordinance**
- **Community engagement**




- **Adult control in specific sites**



ALBO MANAGEMENT PLAN

MOSQUITO CONTROL IN PUBLIC ROAD DRAINS


- census and mapping of public road drains



Quantitative monitoring of *Aedes albopictus*


Emilia-Romagna (4.45 ML residents)

<http://www.zanzaratigreonline.it/>




755 ovitraps in 10 main cities

Emilia-Romagna *Aedes albopictus* egg monitoring



Data source: www.zanzaratigreonline.it

Emilia-Romagna *Aedes albopictus* egg monitoring



Epidemiological equation for VBD

No. bites/person/
day

Vector
competence

Extrinsic cycle
span

Vector
longevity

$$R_0 = \frac{ma^2 * V * P^n}{-\log_e P}$$

Log longevity


from Fine 1981 & Reisen 1989

Aedes albopictus VBD risk estimation

No. eggs/14 days	CHIK A226V	CHIK	DEN	ZIKA
< 100	$R_0 < 1$	$R_0 < 1$	$R_0 < 1$	$R_0 < 1$
101-250	$1 < R_0 < 2$	$R_0 < 1$	$R_0 < 1$	$1 < R_0 < 2$
251-400	$2 < R_0 < 3$	$R_0 < 1$	$R_0 < 1$	$1 < R_0 < 2$
401-700	$3 < R_0 < 5$	$1 < R_0 < 2$	$1 < R_0 < 2$	$2 < R_0 < 3$
701-1000	$5 < R_0 < 7$	$1 < R_0 < 2$	$1 < R_0 < 2$	$3 < R_0 < 5$
1001-1500	$7 < R_0 < 10$	$2 < R_0 < 3$	$2 < R_0 < 3$	$3 < R_0 < 5$
> 1501	$R_0 > 10$	$3 < R_0 < 5$	$3 < R_0 < 5$	$5 < R_0 < 7$

Aedes albopictus monitoring data

Emilia-Romagna 2024




ALBO MANAGEMENT PLAN


Quality control methods on larval treatments in public road drains

- Census and mapping of the public road drains
- PCO/operators must report daily on the treated roads
- Check by GPS tracking system the coverage
- 0.8-1.0 % of the total road drains censused sampled following each treatment
- In small localities a minimum number of 20 road drains must be sampled following each treatment
- Sampling conducted in a pre-defined time window (e.g. 7-14 days post-treatment) with a water net and number of larvae/pupae by species recorded
- Immediate reporting to the Municipality for penalties application

Larvicidal products utilized in public drains treatments in E-R region - 2024



Quality control standard procedure in road drain larval treatment



Opening

Sampling

Cleaning

Counting

EMILIA-ROMAGNA REGIONAL PLAN

Quality control in road drain larval treatment

	2020			2021			2022			2023			2024		
	N. CHEC K	% POS AEDE S	% POS CULE X	N. CHEC K	% POS AEDE S	% POS CULE X	N. CHEC K	% POS AEDE S	% POS CULE X	N. CHEC K	% POS AEDE S	% POS CULE X	N. CHEC K	% POS AEDE S	% POS CULE X
RIMINI	2125	0.2	0.7	3292	2.1	8.3	1881	1.2	2.1	2114	1.8	1.6	1472	2.3	1.8
RICCIONE	1000	1.3	26.5	1000	8.8	35.0	1054	8.5	24.7	2026	1.8	11.1	no data		
RAVENNA	1147	1.1	0.9	1168	0.5	0.5	3233	2.5	0.7	3131	0.9	0.9	2986	2.9	0.4
FORLI'	740	1.6	5.4	727	5.8	5.8	1336	9.7	1.9	1322	2.8	2.1	1284	3.3	1.6
CESENA	508	6.1	4.3	512	7.8	1.4	1236	7.9	1.6	622	4.8	5.5	566	10.4	2.1
IMOLA	3870	3.2	2.6	1491	2.1	1.0	1490	1.1	1.0	1495	1.3	1.5	1484	1.8	1.7
BOLOGNA	1457	4.6	3.2	5892	4.4	6.3	7099	5.5	8.9	7099	4.0	6.7	6652	4.9	7.2
FERRARA	no data			360	4.4	1.9	360	3.1	1.9	984	8.5	4.5	1109	12.2	
MODENA	1659	1.7	1.5	1920	4.5	1.9	3286	7.9	1.8	2494	7.5	3.9	2696	3.4	1.7
REGGIO E.	2651	3.2	2.6	2631	2.2	1.4	2768	3.7	2.4	2912	9.1	5.0	2903	6.4	1.9
PARMA	877	23.6	11.1	709	9.0	8.2	1107	4.2	11.7	766	15.4	7.2	963	3.3	0.4
PIACENZA	no data			no data			672	9.8	6.3	0			420	16.0	6.0
TOTAL	16034			19702			25522			24965			22535		

ALBO MANAGEMENT PLAN

Door-to-door control measures in private areas

- Six larval control treatments in the public road drains (minimum threshold of treated road drains 95%)
- Six door-to-door interventions (larval control, source removal and citizen direct education) (minimum threshold of properties inspected 95%)
- Identification and reporting to the authorities of the non-collaborative citizen
- Introduction of Copepods in large water containers
- Info contact green point (web, phone)
- Socio-economic and environmental impact evaluation

Pilot unit for *Aedes albopictus* sterile male mass-production in Crevalcore – Bologna - Italy



Indicative cost-efficacy *Aedes albopictus* control (Northern Italy)

	€/inhab/ year	% pop reduction	Residual mean no. eggs/ovitrap/14 d
Current IMM	~ 1	~ 30-40	~ 400-600
Intensive DtD	~ 6-8	~ 50-60	~ 200-250
SIT package	~ 9-10	~ 80	~ 100-150

Phone survey on the expenses related to *Aedes albopictus* per E-R family/year

Product	% (N=412)	Cost (€)	N
<i>Insecticide aerosol (domestic)</i>	32.08	6.10	132
<i>Electrical vaporizers</i>	37.86	4.40	156
<i>Outdoor repellents</i>	42.58	4.07	175
<i>Skin repellents</i>	62.53	7.50	257
<i>Mosquito traps</i>	9.95	39.96	41
<i>Larvicides</i>	45.73	3.72	182
<i>Others</i>	1.46	46.29	6
<i>Adulticides</i>	9.71	10.75	40
<i>After bite product</i>	3.88	6.10	16

LIFE Conops project 2013-2018

Laura Vici, Stefano Rivas Morales & Massimo Canali

Dept. of Statistics / Dept. of Food and Agricultural Sciences, University of Bologna

Median annual expense per family

2.4 anti-mosquito products


26.16 – 48.11 Euro

Progetto LIFE Conops 2013-2018

Laura Vici, Stefano Rivas Morales & Massimo Canali

Dept. of Statistics / Dept. of Food and Agricultural Sciences, University of Bologna

Any questions or comments ?



Overview du contrôle de la transmission des arboviroses liées à Aedes

Protéger les communautés avec Wolbachia - Indonésie

par Edwige Rancès

Protéger les communautés avec ***Wolbachia***

INDONÉSIE

Edwige Rancès



Qui sommes nous?

Organisation à but non lucratif
appartenant à l'Université Monash,
Australie.

Bureaux en Australie et en France

Équipe internationale et multidisciplinaire




Construisons avec les gouvernements

Des partenariats stratégiques fondés sur la co-crédation et l'intégration à long terme.

C'est ainsi que nous protégeons davantage de personnes, de manière plus durable.



Wolbachia dans une cellule d'insecte



Plus de 50% des populations d'insectes portent *Wolbachia*

- **Wolbachia**, outil de lutte biologique contre :



Réduction de l'incidence de la dengue après l'introgression de *Wolbachia* dans les populations naturelles d'*Aedes aegypti* : essai randomisé, séries chronologiques.





Les résultats des analyses socio-économiques (coût/efficacité) prédisent *Wolbachia* comme étant une solution efficace.

En moyenne, chaque BRL investi **devrait rapporter 278,73 BRL en bénéfices économiques.**

Zimmermann et al., 2024 The Lancet Regional Health Americas

En moyenne, chaque dollar investi **devrait rapporter 1,69 \$ en bénéfices économiques sur 20 ans**

Turner HC, et al. PLoS Negl Trop Dis 2023.

Mexico

El Salvador Honduras

Colombia

Brazil

Senegal

Sri Lanka

Laos
Vietnam

Indonesia
Timor Leste

Australia

Kiribati
Vanuatu

Fiji

New Caledonia

Les avantages économiques sur 10 ans dépassent 5,00 \$ US pour chaque dollar investi

Shepard et al., 2025

Rembourse 2 à 3 fois l'investissement sur 10 ans grâce aux économies réalisées sur les frais médicaux et la perte de productivité

Brady OJ, et al. BMC Medicine 2020.

Une seule épidémie de dengue (coût de santé et mesure de lutte antivectorielle) coûte 2 fois l'investissement réalisé.
Une épidémie de dengue a coûté 13 millions d'euros.

Une intervention *Wolbachia* a coûté 6,5 millions d'euros.

DASS New Caledonia Government.



Depuis 2011 (premiers essais opérationnels), nous travaillons en partenariat avec 16 pays





543km²




1.8 million de personnes protégées



**World
Mosquito
Program**



UNIVERSITAS GADJAH MADA



2014-2015

Pilotes périurbains

2016-2019

Étude quasi-expérimentale (2006-2019)

Déploiement de *Wolbachia* dans la zone expérimentale (2016)

2017-2020

Essai clinique randomisé par grappes

Déploiement de *Wolbachia* dans les clusters expérimentaux (2017)

2020-2021

Déploiement de *Wolbachia* dans les zones/clusters témoins

2021-2022

Déploiement de *Wolbachia* dans la province de Sleman


2022-2023

Déploiement de *Wolbachia* dans la province de Bantul

2023-2025


Étude au long terme post-intervention






2016-2019 Étude quasi-expérimentale

- *Objectif principal* : Évaluer l'impact du déploiement de *Wolbachia* sur l'incidence de la dengue.
- *Intervention* : Introgression de *Wolbachia* dans les populations d'*Aedes aegypti* (Aout 2016 – Mars 2017)
 - Zone intervention : 65 000 habitants, 4,9 km²
 - Zone témoin : 34 000 habitants, 3.1 km²





76% réduction (95% CI: 60%, 86%) de l'incidence de la dengue
 Étude de séries chronologiques de cas de dengue notifiés (2006-2019)






2016-2020 Étude expérimentale

- Essai clinique randomisé par grappes (clusters)
- *Objectif principal* : Évaluer l'efficacité du déploiement de *Wolbachia* sur la réduction de l'incidence de cas de dengue confirmés virologiquement, sur une population âgée de 3 à 45 ans.
- *Objectif secondaire* : Évaluer l'efficacité du déploiement de *Wolbachia* sur la réduction de l'incidence de cas de chikungunya et de zika confirmés virologiquement, sur une population âgée de 3 à 45 ans.
- *Intervention* : Introgression de *Wolbachia* dans les populations d'*Aedes aegypti* (Mars 2017-Décembre 2017)
 - Zone intervention : 12 clusters, approx 1 km²
 - Zone témoin : 12 clusters, approx 1 km²
- Nombre d'habitants : 311 700






● Participants (cumulative)

✚ Puskesmas




Au total, 385 cas de dengue ont été inclus (dont 328 sérotypés) 67 dans le groupe d'intervention ; 318 dans le groupe non traité


***Wolbachia* perturbe le regroupement spatio-temporel des cas de dengue**



Sous-ensemble de 160 cas de dengue regroupés dans l'espace et le temps




L'introggression de *Wolbachia* n'a pas d'impact sur la prévalence d'*Aedes aegypti*



4 BG traps/km² clusters intervention ; 5 BG traps/km² clusters témoins avant l'introgession de *Wolbachia*


14 BG traps/km² clusters intervention ; 15 BG traps/km² clusters témoins pendant et après l'introgession de *Wolbachia*

L'introgession de *Wolbachia* n'a pas d'impact sur la prévalence d'*Aedes albopictus*




Cypermethrin in 2016-2018
Malathion in 2019-2020

Diminution importante des épisodes de pulvérisation d'insecticides




2023- 2025 Étude au long terme post-intervention sur Yogyakarta


- Étude de séries chronologiques de cas de dengue sévère notifiés (2006-2025)
- *Objectif principal* : Évaluer l'impact du déploiement de *Wolbachia* sur le taux d'hospitalisation de cas de dengue.
- Zone intervention : 400 000 habitants ; 32 km²



% d'*Aedes aegypti* positif pour *Wolbachia* sur une échelle de temps de 2017 à 2025 par village / sous-district



89% des *Aedes aegypti* collectés portent *Wolbachia* en 2024



Estimation d'une réduction de 78% de l'incidence de la dengue (cas hospitalisés)
Etude de séries chronologiques (2006-2025)



World
Mosquito
Program



Yayasan
TAHIYA




UNIVERSITAS
GADJAH MADA

UC Berkeley



MONASH
University

Extra slides




Wolbachia est transmise de façon **maternelle** et peut manipuler la reproduction de ses hôtes afin de favoriser son introgression dans les populations naïves (**incompatibilité cytoplasmique**).



Quelle est la méthode *Wolbachia* développée par WMP?

- 1** En laboratoire, introgression de *Wolbachia* dans une population autochtone d'*Aedes aegypti*, suivie d'une série de contrôles qualité (notamment l'interférence virale), avant de passer en production de masse et lâchers.




- 2** *Wolbachia-Aedes aegypti* sont lâchés durant 3 à 6 mois.



- 3** Les moustiques lâchés s'accouplent avec les populations sauvages d'*Aedes aegypti* et passent *Wolbachia*.



- 4** Lorsque les lâchers sont terminés, *Wolbachia* se répand dans la quasi-totalité des populations d'*Aedes aegypti* sauvages et se maintient.



- 5** Les communautés sont protégées par *Wolbachia* durablement et sans la nécessité de faire de nouveaux lâchers.






Moustiques
avec *Wolbachia*



Moustiques sans
Wolbachia







- High and stable *Wolbachia* frequencies (average 93%) in intervention clusters throughout trial
- Some 'contamination' of untreated clusters after 1 year into clinical enrolment

Nouvelles méthodes de LAV : techniques en phase terrain

Wolbachia : stratégies de remplacement et de suppression

par Nicolas Pocquet

***Wolbachia* :**

Stratégies de remplacement et de suppression

Nicolas POCQUET (PhD, HDR)


Responsable de l'unité d'entomologie médicale
Institut Pasteur de Nouvelle-Calédonie
npocquet@pasteur.nc




Contexte général

La dengue

- Dans le monde
 - Nombre de cas multiplié par 10 en 20 ans (2000 – 2019)
 - Répartition mondiale
 - **> 5 millions de cas en 2023 → 5000 décès**
- Dans les pays et territoires insulaires du Pacifique
 - 9 pays touchés en 2023 (n = 21)
 - **> 13 000 cas en 2023** (28 % d'augmentation vs 2022)
- Au delà de l'impact humain...
 - Impact économique
 - Mondial 1975 – 2022 → **76,5 milliards US\$**
(Roiz et al., 2024)
 - Epidémie Nouvelle-Calédonie 2013 → **13 millions €**
(estimation DASS-NC)



Carte des pays ayant signalé des cas de dengue autochtones, nov. 2022 à nov. 2023
(www.who.int ; WHO 2023)



Evolution et coût annuel (en millions US\$) de la dengue, du chikungunya et du Zika
(Roiz et al., 2024)


Limites de la LAV : l'exemple de la Nouvelle-Calédonie

La Dengue en NC


- Un seul vecteur de DENV : *Aedes aegypti*

Efficacité limitée de la LAV conventionnelle

- **Communication / mobilisation sociale :**
 - Efficace sur les indices entomologiques
 - Pas d'impact visible sur les épidémies
- **Insecticides :**
 - Traitements adulticides peu efficaces (e.g., ULV)
 - Augmentation de la **résistance à la deltaméthrine**
 - Pas d'insecticide alternatif
 - Réticence de la population



Moyennes du nombre de larves et nymphes retrouvées par maison à Nouméa entre 2000 et 2017




Evolution de la résistance à la deltaméthrine à Nouméa entre 2003 et 2018
(IPNC, données non publiées)

Nouvelles techniques de lutte basées sur *Wolbachia*


Technique de l'Insecte Incompatible vs Remplacement

- Propriétés de *Wolbachia*
 - Incompatibilité Cytoplasmique (IC)
 - Blocage de la réplication virale (DENV, ZIKV, CHIKV, YFV)

a Suppression of mosquito population Cytoplasmic incompatibility



b Modification of mosquito population



Comparaison des techniques basées sur l'utilisation de *Wolbachia* – (a) Technique de l'Insecte Incompatible ou TII, (b) Remplacement ou modification de la population (Flores & O'Neill, 2018)

Nouvelles techniques de lutte basées sur *Wolbachia*

Technique de l'Insecte Incompatible (TII)

Avantages

- Utilisable sur plusieurs vecteurs de DENV :
 - *Ae. aegypti*
 - Espèces du groupe Scutellaris (e.g., *Ae. albopictus*)
 - Autres vecteurs d'arbovirus (e.g., *Culex spp.*)
- Réduction de la nuisance
- Réduction de la transmission (e.g., Singapour)


Inconvénients

- Production de masse et lâchers conséquents (comparable TIS)
- Importance du sexage pour éviter l'installation de *Wolbachia* :
 - TII + irradiation légère (stérilisation des femelles)
 - TII + nouvelles techniques de tris des nymphes
- Non durable en milieu ouvert (i.e., Singapour)

Dans les territoires ultra-marins

- Expérimentation de terrain depuis 2015 sur *Ae. polynesiensis* en **Polynésie Française**
- Très bons résultats sur un atoll de PF, en association à des traitements larvicides (www.ilm.pf, 2025)

a Suppression of mosquito population
Cytoplasmic incompatibility



Technique de l'Insecte Incompatible ou TII
(Flores & O'Neill, 2018)

Nouvelles techniques de lutte basées sur *Wolbachia*

Remplacement de population

Avantages

- Production de masse et lâchers réduits
- Pas de sexage nécessaire
- Durable une fois *Wolbachia* établie
- Réduction importante de la transmission (*i.e.*, Indonésie : OR = 0,23)


Inconvénients

- Uniquement utilisable sur *Ae. aegypti*
- Pas de réduction de la nuisance

Dans les territoires ultra-marins

- Implémentée depuis 2018 en **Nouvelle-Calédonie**

b Modification of mosquito population



*Remplacement ou modification de la population
(Flores & O'Neill, 2018)*

Le World Mosquito Program

- Technique de remplacement basée sur *Wolbachia*
- Programme à but non lucratif
- Porté par la Monash University
- Implémenté dans 14 pays (depuis 2011 en Australie)

Cadre du projet en NC

- **Financeurs** : Gouv NC, Université de Monash (WMP), IPNC, Villes de Nouméa, Mont-Dore, Dumbéa, Païta, Fonds Pacifique, Province Sud
- **Collaborateurs** : WMP, IPNC, DASS-NC, Mairies de Nouméa, Mont-Dore, Dumbéa et Païta
- **PI** : Nadège ROSSI
- **Budget global** : 6,7 M€ (subvention : 3,5 M€)
- **Période** : 2018 – 2024




WMP-NC :

Souche NC-wMel


Création de la souche

- Souche **NC-wMel** générée à partir de la souche wMel australienne
 - 6 étapes de Backcross
 - Mâles **WT** x Femelles wMel
- Obtenir une souche génétiquement proche des moustiques **WT** (98,5 %)
- Maintien de la souche **NC-wMel**
- Contrôle de la fréquence de *Wolbachia* à chaque génération



Tests de Fitness

- **Transmission maternelle**
 - Croisements ♂ WT x ♀ NC-wMel
 - 96 % de transmission de *Wolbachia* à la descendance
- **Incompatibilité Cytoplasmique**
 - Croisements ♂ NC-wMel x ♀ WT
 - 0 % d'éclosion des œufs : IC complète
- **Fécondité et Fertilité**
 - Croisements ♂ NC-wMel x ♀ NC-wMel
 - 22 œufs par femelle (43 pour la souche WT)
 - 53 % d'éclosion des œufs (77 % pour la souche WT)
- **Résistance aux insecticides**
 - Paramètre important pour les lâchers
 - Forte résistance de la souche NC-wMel ($RR_{50} = 42$)
 - Niveau de résistance équivalent aux WT



Tests doses réponses à la deltaméthrine sur les souches bora (sensible) WT et NC-wMel (résistantes) (Pocquet et al., 2021)

WMP-NC :


Blocage de l'infection

Repas infectieux

- Virus ayant circulés en NC
 - DENV-2
 - CHIKV
 - ZIKV

→ Réduction significative du taux d'infection pour la souche **NC-wMel**

→ Pas de virus détecté dans la salive




Taux d'infection et efficacités de transmission pour les souches d'Ae. aegypti NC-WT et NC-wMel exposées à des repas de sang infectés par du DENV-2, du CHIKV ou du ZIKV. (Pocquet et al., 2021)

Injections intrathoraciques

- Virus de référence
- DENV-1 à 4

→ Réduction significative du taux d'infection pour la souche **NC-wMel**, comparable à la souche **AUS-wMel**

→ Réduction significative du nombre de copies de DENV chez les moustiques **NC-wMel** infectés vs **NC-WT** infectés



Taux d'infection et nombre de copie d'ARN viral pour les souches d'*Ae. aegypti* WT et wMel (NC et Australie) exposées à des micro-injections des 4 sérotypes de DENV. (Pocquet et al., 2021)

Autorisation de lâchers

- NC : autonome sur les questions d'environnement et de santé
- Comité d'éthique de Nouvelle-Calédonie
- Autorisation du gouvernement de la Nouvelle-Calédonie

Acceptation de la population

- Campagne de communication et engagement communautaire
- Evaluation de l'acceptation du projet par la population
- Constitution d'un groupe de référence communautaire
- Sondage de la population :
 - 94% de personnes favorables à Nouméa
 - 91% et 93% à Dumbéa et Mont Dore (2 phases)
 - 94% à Païta



*Campagne de communication WMP-NC Nouméa
(WMP-NC, 2018)*

WMP-NC :

Déploiement

Phase I (Nouméa)

- Août 2019 – Juillet 2021
- 2800 points de lâchers / 230 BG
- 12 M de moustiques relâchés / 26 à 63 semaines de lâchers

Phase II (Dumbéa / Mt Dore)


- Janvier 2021 – Juillet 2022
- 2000 points de lâchers / 120 BG
- 4 M de moustiques / 24 semaines

Phase III (Dumbéa / Mt Dore)

- Novembre 2022 – Juin 2023
- 1400 points de lâchers / 50 BG
- 4 M de moustiques / 22 semaines

Phase IV (Païta)

- Novembre 2023 – Mai 2024
- 1400 points de lâchers / 93 BG
- 4 M de moustiques / 24 semaines



WMP-NC :

Résultats entomologiques

Phase I (Nouméa)

- 87 % de moustiques porteurs de *Wolbachia* 33 mois post-lâchers

Phase II (Dumbéa / Mt Dore)


- 92 % de moustiques porteurs de *Wolbachia* 21 mois post-lâchers

Phase III (Dumbéa / Mt Dore)

- 78 % de moustiques porteurs de *Wolbachia* 10 mois post-lâchers

Phase IV (Païta)

- 80 % de moustiques porteurs de *Wolbachia* 6 mois post-lâchers



Suivi épidémiologique

- Suivi des cas de dengue par les autorités de santé (**DASS-NC**)
 - Pas d'épidémie depuis 2019
 - Faible transmission autour des cas importés

Suivi long terme en cours


- Suivi entomologique une fois par an
- Suivi épidémiologique en continu
- Projets de recherche en parallèle

*Suivi mensuel des cas de dengue
autochtones et importés en NC
entre 2008 et 2025*

(DASS-NC, données non publiées)


Suivi mensuel des cas de dengue déclarés en Nouvelle-Calédonie

01/01/2008 au 17/02/2025



Suivi mensuel des cas de dengue importés déclarés en Nouvelle-Calédonie

01/01/2008 au 17/02/2025



Projet PYRWOL (IPP / IPNC)


- Bourse de thèse ANRS-MIE / Arbo-France
- Populations de terrain échantillonnées en 2024 :
 - Nouméa : *Ae. aegypti* **wMel**
 - Ouvéa : *Ae. aegypti* **WT**
- Infections expérimentales au DENV2



Dupuis B.

Résultats

- Infection
 - **WT** : 88 % à 14 dpi (maximum à 100 %)
 - **wMel** : 16 % à 14 dpi (maximum à 17 %)
- Efficacité de transmission
 - **WT** :
 - Premiers moustiques infectants à 14 dpi
 - $EIP_{50} = 18$ jours (IC_{95} : 13 – 21)
 - 64 % de moustiques infectants au maximum
 - **wMel** :
 - Premiers moustiques infectants à 28 dpi
 - EIP_{50} non calculable
 - 6 % de moustiques infectants au maximum



Taux d'infection et efficacités de transmission du DENV2 par des populations d'*Ae. aegypti* prélevées en NC en 2024 (Dupuis et al., données non publiées)

Conclusion

Contrôle des vecteurs et des arboviroses :

- Traitements conventionnels (ULV et thermo-nébulisation) encore très présents
- Efficacité entomologique faible, voir inexistante sur populations résistantes
- Efficacité épidémiologique peu évaluée

Wolbachia : outil d'intérêt en LAV

- Recherches ET application de la recherche en cours :
 - **TII :**
 - 8 essais de terrain de TII ou TII + TIS publiés (7 entre 2019 et 2024)
 - *Ae. albopictus* (2) et *Ae. aegypti* (6)
(Revue dans Rahul *et al.*, 2024)
 - **Remplacement :**
 - Implémenté dans 14 pays sur *Ae. aegypti* (WMP)
 - Technique d'intérêt pour la Guyane, la Guadeloupe et la Martinique
(i.e., *Ae. aegypti* seul vecteur de DENV)



Merci pour votre attention

Nouvelles méthodes de LAV : techniques en phase terrain

TIS classique et TIS boostée

par Jeremy Bouyer


Journées scientifiques :
Moyens innovants de lutte contre les vecteurs d'arbovirus.
Institut Pasteur, Paris

TIS classique et TIS boostée

Jérémy Bouyer
23 Juin 2025



Sterile Insect Technique (SIT)




<http://www.naweb.iaea.org/nafa/resources-nafa/SIT-Mosquitoes-LR.mp4>



Joint FAO/IAEA Centre
Nuclear Techniques in Food and Agriculture

Sterile Insect Technique (SIT): impact of mating harassment...



nature communications



Article

<https://doi.org/10.1038/s41467-024-46268-x>

Mating harassment may boost the effectiveness of the sterile insect technique for *Aedes* mosquitoes




Received: 21 July 2023

Accepted: 20 February 2024

Dongjing Zhang^{1,14}, Hamidou Maiga^{2,3,14}, Yongjun Li^{4,5,14},
Mame Thierno Bakhom^{3,6,14}, Gang Wang^{1,14}, Yan Sun¹, David Damien⁷,
Wadaka Mama², Nanwintoun Séverin Bimilé Somda^{2,8}, Thomas Wallner²,
artina², Simran Singh Kotla², Hanano Yamada²,
Jiatian Guo¹, Qingdeng Feng¹, Junyan Zhang¹,
Wenjie Pan¹⁰, Yu Wu¹, Xiaoying Zheng¹,
Marc J. B. Vreysen² & Jérémy Bouyer^{2,12,13,14} ✉



Boosted SIT




Pyriproxyfen

(Bouyer & Lefrançois, 2014)



SIT & IIT-SIT & Boosted SIT: ONGOING AND PAST TRIALS (2024)



Bouyer *Infectious Diseases of Poverty* (2024) 13:68
<https://doi.org/10.1186/s40249-024-01242-z>


Infectious Diseases of Poverty

COMMENTARY

Open Access

Current status of the sterile insect technique for the suppression of mosquito populations on a global scale

Jérémy Bouyer^{1,2,3*}



Legend

Strategic approach
in testing sites

- IIT-SIT (4)
- SIT (32)
- ◆ Boosted SIT (3)

SIT & IIT-SIT & Boosted SIT: ONGOING AND PAST TRIALS (2024)

Bouyer *Infectious Diseases of Poverty* (2024) 13:68
<https://doi.org/10.1186/s40249-024-01242-z>

Infectious Diseases of Poverty

COMMENTARY

Open Access

Current status of the sterile insect technique for the suppression of mosquito populations on a global scale



Jérémy Bouyer^{1,2,3*}

Phase	I	II	III	IV	Total
2019	21	11	2	0	34
2024	12	22	4	1	38
Increase rate	-43%	100%	50%	100%	NA

IIT-SIT: Phase IV in Singapore

Effectiveness of *Wolbachia*-mediated sterility coupled with sterile insect technique to suppress adult *Aedes aegypti* populations in Singapore: a synthetic control study

Somya Bansal*, Jue Tao Lim*, Chee-Seng Chong*, Borame Dickens*, Youming Ng, Lu Deng, Caleb Lee, LiYun Tan, Evdoxia G Kakani, Yanni Yoong, David Du Yu, Grace Chain, Pei Ma, Shuzhen Sim, Lee Ching Ng, Cheong Huat Tan
Lancet Planet Health 2024;
8: e617–28



“We found that *Wolbachia*-based IIT-SIT reduced wild-type female *A aegypti* populations by a mean of 62·01% (95% CI 60·68 to 63·26) by 3 months, 78·40% (77·56 to 79·18) by 6 months, and **91·32% (90·95 to 91·66) by at least 18 months** of releases.”


“Our study comprised an at-risk population of 607 872 individuals living in intervention sites and 3 894 544 individuals living in control sites. Interventions demonstrated up to **77·28% (121/156, 95% CI 75·81–78·58)** intervention efficacy despite incomplete coverage across all towns until EW 26, 2022.”

THE STRAITS TIMES

SINGAPORE

Project Wolbachia to expand to five more sites in S'pore to lower dengue risk









SIT in Reunion island




- Phase II trial in Sainte Marie 20 ha
- Mass-production module established (CRVOI)
- 300.000 sterile males produced / released per w in 2021
- Excellent quality of the sterile males
- Max 50% of induced sterility (target area too small & not isolated) / no reduction of adult density
- Very good acceptance by the beneficiary population



Beware of (over)-compensation!




Value of SIT in AW-IPM strategies



Joint FAO/IAEA Centre
Nuclear Techniques in Food and Agriculture

Model predictions



Better suppression & resilience than SIT

Boosted SIT phase II trials

Aedes aegypti (Reunion)

St Joseph, Réunion 10ha




>90% reduction of adult *Aedes aegypti* in 3 months

~60% reduction of adult *Aedes albopictus* & eggs in 2 months

Aedes albopictus (Spain)

Vilavella, 35ha




50-95% reduction of adult *Aedes albopictus*

25-60% reduction of eggs *Aedes albopictus*

14-26% increased suppression in comparison to SIT


Polinyà de Xúquer, 22ha





50-95% reduction of adult *Aedes albopictus*

25-60% reduction of eggs *Aedes albopictus*

Boosted SIT phase II trial in Greece / intermittent releases



Trapping date
Release date



<8% egg population reduction in the treated site compared to the control site

Reduction of adults started 1 month after the first cycle of release (40%), reached 85-93% during the second cycle and for one month

Main results of boosted SIT trials in Europe (phase II)



- More efficient than SIT, but not as much as predicted in Spain (large larval habitats?)
- Partially protects against “eggs immigration”
- Prevents compensation & overcompensation
- Persistent effect
- Perfect combination with SIT
- No observed environmental impact (bees & chironomides in Reunion)
- Excellent acceptance by beneficiary populations

Perspectives



pTIS

Phase III trial in Reunion 2025-26



Saint Joseph



OBJECTIVES

- ➔ Sequential upscale 50->175ha conditional to success & acceptability
- ➔ Efficiency on *Ae. albopictus* & *Ae. aegypti*
- ➔ Environmental impact (water, chironomides & other NTOs, bees)
- ➔ Impact on vector-host contact & dengue incidence
- ➔ Economical model with transfer to a local start-up




Suivi environnemental


4 ruchers avec 10 ruches par site avec suivi de la santé des ruches (méthode Coleval) et mesure des résidus de pyriproxifène dans la cire, le pollen, le miel et les abeilles tous les 3 mois (sous-traitance *CETA*)

Suivi de 15 gîtes larvaires par zone (10 naturels et 5 anthropiques):

- dosage du pyriproxifène dans l'eau des gîtes larvaires
- Faune non-cible de ces gîtes caractérisée tous les mois (taxonomie & eDNA)
- Faune non-cible comptabilisé dans les pièges à adultes (bycatch)




Localisation des ruchers sentinelles et aires de récolte des abeilles



Creation of a start-up spin-off from Cirad in Sept. 2023

Mosquito control service
(450-2000€/ha/y)

- Corporates (hotels, islands)
- Municipalities (schools, hospitals)



Demande d'opérationnalisation accélérée par 4 mairies de la Réunion suite à l'épidémie de chikungunya

Etude de faisabilité en cours pour une usine avec une capacité de 2-2,5 millions de mâles stériles à St Joseph

st JOSEPH
LA RÉUNION 974
LE SUD SOVAJ'



Utilisation hétérospécifique de la TIS boostée pour prévenir les invasions

Trends in Parasitology


Available online 21 June 2025

In Press, Corrected Proof [What's this?](#)

Forum

Preventing the establishment of invasive exotic mosquitoes

Jérémy Bouyer^{1,2,3}, Diana Iyaloo⁴, Thierry Baldet^{1,2}



Merci!




Joint FAO/IAEA Centre
Nuclear Techniques in Food and Agriculture

Nouvelles méthodes de LAV : techniques en phase terrain

Genetically modified mosquitoes for disease control : from conventional transgenesis... to gene drive ?


par Eric Marois

GM mosquitoes for disease control : from conventional transgenesis... to gene drive ?




Oxitec : RIDL and fs-RIDL technologies

Release of Insects carrying a female-specific Dominant Lethal gene



Mass-produce
with tetracycline



→ Release eggs

→ females die,
sterile males
mate with wild
females




→ Daughters die ;
sons transmit gene
again to 50% progeny



→ Population
suppression

Oxitec interventions




RIDL and fs-RIDL Oxitec technologies

Advantages & drawbacks

- Species-specific
- Self-dissemination
- Circumvents insecticide resistance
- No need for irradiation $\neq$ SIT
- No need for sex separation in fs-RIDL $\neq$ SIT
- Self-limiting
- Costly, complex logistics (less with fs-RIDL)
 - field trials rather than full-blown interventions
- Small spatial scale
- Not long-lasting
- Public reluctance on GM, depending on countries

SIT: Transgenes to separate males / females




- COPAS sorting of young larvae
- irradiation
- Sterile Insect Technique

(Lutrat et al., 2022, 2023)

Nix-expressing transgenes are sufficient to convert females into males




Célia Lutrat
et al. Comm Biology
2022, 2023




Plan A (most radical) : Maleness gene drives

Dock genes encoding the CRISPR-Cas9 system into synthetic M loci :




In spermatogonia, chromosome pair with heterozygous *Nix* transgene:




**A pseudomale siring only pseudosons
= self-sustaining, masculinizing gene drive !**

Plan B (less radical): M-linked editors

Dock genes encoding the CRISPR-Cas9 system into synthetic M loci :




→ YLE (Y-linked editors) (Burt & Deredec, 2018) → MLE:





→ males whose daughters are non-flying or sterile

Mathematical modeling : not self-sustaining, but more efficient than SIT or fsRIDL




Now get CRISPR/Cas9 to work efficiently in *albopictus*

- clone better germline promoters to express Cas9
- Find the best expression system for multiplex gRNAs




Technology development: test promoters to optimize Cas9 and gRNA expression

candidate promoter-Cas9 lines



line expressing 3 gRNAs targeting yellow

X




→ F1 : mutagenesis of *yellow* in the germ line
x *yellow* mutant

→ F2: count the proportion of *yellow*




→ Optimal promoter for gRNAs

→ Optimal promoter for Cas9



masculinizing gene drive





Célia Lutrat



Julie Fimeyer



Marie Aubry



Nikola Djordjevic



Victor Redon



Mateus Berni



Lucie Souriac



Roénick P. Olmo



Amandine Gautier



Stéphanie Blandin





Phi Papathanos



Doron Zaada



Jérémy Bouyer



anr®

Université

de Strasbourg




Inserm




La science pour la santé
From science to health

Anopheles gambiae dual *Saglin* / *Lipophorin* gene drive

(Green et al., eLife 2024)



Rapid invasion by both transgenes



Nouvelles méthodes de LAV : techniques en phase terrain

Un redoutable envahisseur arrive dans les Antilles Françaises

par Anubis Vega Rua

A fearsome invader arrives in the French West Indies



Anubis Vega-Rúa
Vector-Borne Diseases Laboratory
Institut Pasteur de Guadeloupe

Colloque Arbofrance, June 23th 2025



The French Territories of the Americas





The French Territories of the Americas



The French Territories of the Americas



The French Territories of the Americas



Leon et al, *In preparation*


The French Territories of the Americas



Espèce (*Aedes* spp.)

- *Aedes aegypti*
- *Aedes albopictus*
- *Aedes polynesiensis*

The French Territories of the Americas



Espèce (*Aedes* spp.)

- *Aedes aegypti*
- *Aedes albopictus*
- *Aedes polynesiensis*

Tropical Medicine and International Health

doi:10.1111/j.1365-3156.2010.02613.x

VOLUME 16 NO 1 PP 134-139 JANUARY 2011

Viewpoint

High susceptibility to Chikungunya virus of *Aedes aegypti* from the French West Indies and French Guiana

Romain Girod¹, Pascal Gaborit¹, Laurence Marrama², Manuel Etienne³, Cédric Ramdini⁴, Ignace Rakotoarivony², Christelle Dollin², Romuald Carinci¹, Jean Issaly¹, Isabelle Dusfour¹, Joël Gustave⁴, Marie-Michelle Yp-Tcha³, Andre Yébakima³, Anna-Bella Failloux³ and Marie Vazeille⁵

Emerging Microbes & Infections

2019, VOL 8

<https://doi.org/10.1080/22221751.2019.1615849>




Taylor & Francis
Taylor & Francis Group

OPEN ACCESS

Transmission potential of African, Asian and American Zika virus strains by *Aedes aegypti* and *Culex quinquefasciatus* from Guadeloupe (French West Indies)



Lyza Hery*, Antoine Boullis*, Christelle Delannay and Anubis Vega-Rúa



bio
RENDER


INSTITUT
PASTEUR
de la Guadeloupe

On September 30th, 2024...



Vector control agents collect mosquito larvae from a floor siphon in Saint Barthélemy

On September 30th, 2024...




Surprise!


Strong suspicion of *Aedes albopictus*



On September 30th, 2024...



Strong suspicion of *Aedes albopictus*



Mission : October 14th-16th, 2024



Objectives:

- Confirm *Ae. albopictus* identification morphologically
- Train the local agents on *Ae. albopictus* identification of diverse mosquito sampling techniques
- Event entomological monitoring
- Recover specimens to confirm identification by other approaches:
 - Matrix Assisted Laser Desorption Ionization - Time-of-Flight Mass Spectrometry (MALDI-TOF MS)
 - Cytochrome oxidase 1 (*cox1*) barcoding

Mission : October 14th-16th, 2024



Objectives:

- Confirm *Ae. albopictus* identification morphologically
- Train the local agents on *Ae. albopictus* identification of diverse mosquito sampling techniques
- Event entomological monitoring
- Recover specimens to confirm identification by other approaches:
 - Matrix Assisted Laser Desorption Ionization - Time-of-Flight Mass Spectrometry (MALDI-TOF MS)
 - Cytochrome oxidase 1 (*cox1*) barcoding



Aedes scutellaris



Aedes polynesiensis




MALDI-TOF MS approach confirmed *Ae. albopictus* ID


Legs



MS
biotyping



Mosquito species
identification



Aedes albopictus

Aedes polynesiensis

Aedes malayensis

Aedes pseudoscutellaris

Aedes futunae


Aedes scutellaris



Ramdini *et al.* in revision




Cytochrome oxidase 1 barcoding confirmed *Ae. albopictus* ID



Event entomological surveillance: October 14th -16th, 2024.



Event entomological surveillance: October 14th -16th, 2024.




Aedes albopictus breeding sites identified, Saint Barthélemy

Breeding site	Coordinates	Species identified*
Floor Siphon (post office)	17°54'20"N, 62°49'28"W	<i>Ae. aegypti</i> ; <i>Ae. albopictus</i>
Tire	17°54'16"N, 62°49'27"W	<i>Ae. aegypti</i> (6); <i>Ae. albopictus</i> (2; males)
Watering can	17°54'15"N, 62°49'32"W	<i>Ae. albopictus</i>
Flowerpot dish	17°54'21"N, 62°49'29"W	<i>Ae. aegypti</i> (10); <i>Ae. albopictus</i> (1; female)
Small container	17°54'24"N, 62°49'15"W	<i>Ae. albopictus</i>

*The number of specimens (when available) is given in parenthesis.

Event entomological surveillance: October 14th -16th, 2024.



Aedes albopictus breeding sites identified, Saint Barthélemy


Breeding site	Coordinates	Species identified*
Floor Siphon (post office)	17°54'20"N, 62°49'28"W	<i>Ae. aegypti</i> ; <i>Ae. albopictus</i>
Tire	17°54'16"N, 62°49'27"W	<i>Ae. aegypti</i> (6); <i>Ae. albopictus</i> (2; males)
Watering can	17°54'15"N, 62°49'32"W	<i>Ae. albopictus</i>
Flowerpot dish	17°54'21"N, 62°49'29"W	<i>Ae. aegypti</i> (10); <i>Ae. albopictus</i> (1; female)
Small container	17°54'24"N, 62°49'15"W	<i>Ae. albopictus</i>

*The number of specimens (when available) is given in parenthesis.

Mixed breeding sites

Event entomological surveillance: Late October-November, 2024.

Near the airport!



Event entomological surveillance: Late October-November, 2024.




December: Insect control campaign (*Bti*, breeding sites reduction, **deltamethrin treatments** in areas surrounding positive breeding sites)


Entomological surveillance update: June, 2025.




Ae. albopictus in the Caribbean: decades of colonisation



Ae. albopictus in Saint Barthélemy : the beginning of a new colonization wave?



Saint Barthélemy aerial connections



<https://www.flightconnections.com>

What to do in terms of vector control and surveillance?

Currently ongoing:

- Weekly, entomological surveillance is conducted in Saint Barthélemy by 2 agents (30 ovitraps, larval prospections)
- Occasional reinforcements with agents from Saint Martin (3) and/or Martinique.

Needs:

- Reinforcement of local agents (to increase the scope of vector surveillance and control actions).
- Saint Martin and Guadeloupe: Reinforce staff for entomological surveillance, particularly at ports of entry

What to do in terms of research

Currently ongoing:



- Weekly, entomological surveillance is conducted in Saint Barthélemy by 2 agents (30 ovitraps, larval prospections)
- Occasional reinforcements with agents from Saint Martin (3) and/or Martinique.


Needs:

- Reinforcement of local agents (to increase the scope of vector surveillance and control actions).
- Saint Martin and Guadeloupe: Reinforce staff for entomological surveillance, particularly at ports of entry

Research:


- Strengthen collaboration between ARS agents and research institutions to go deeper into surveillance data analysis and provide insights about vector ecology
- Characterize vector resistance and identify its importation origin
- Entomo-virological surveillance
- Vector competence

Acknowledgements



Thank you for your attention

Confirmation par MALDI-TOF MS



Mix	Detected Species	Log(Score)
1	Aedes albopictusfemelle-ref-4 22-IPM	2.050
2	Aedes albopictus Leqs CMR	2.020
3	Aedes albopictus Leqs MRS	1.940
4	Aedes albopictus-ref-2 20-IPM	1.890
5	Aedes albopictus Leqs DOU	1.860
6	Aedes albopictus-Ref-CA-074-IPNC	1.830
7	Aedes albopictus-ref-Fiji-021-IPNC	1.800
8	Aedes albopictus-Ref-CA-073-IPNC	1.800
9	Aedes albopictus-ref-Fiji-021-IPNC	1.790
10	Aedes albopictusfemelle-ref-IPM1 22-IPM	1.770

Mix	Detected Species	Log(Score)
1	Aedes albopictus Thorax MRS	2.050
2	Aedes albopictus Thorax CMR	1.970
3	Aedes albopictus-ref-Fiji-021-IPNC	1.510
4	Aedes albopictus-Ref-Fiji-021-IPNC	1.490
5	Aedes albopictus Leqs MRS	1.470
6	Aedes albopictus-ref-Fiji-021-IPNC	1.470
7	Aedes pseudoscutellaris-Ref-Fiji-314-IPNC	1.390
8	Aedes albopictus-Ref-LA-030-IPNC	1.370
9	Aedes pseudoscutellaris-Ref-Fiji-163-IPNC	1.310
10	Aedes albopictus-Ref-Fiji-018-IPNC	1.280

Range	Description	Symbols	Color
2.300 ... 3.000	highly probable species identification	(+++)	green
2.000 ... 2.299	secure genus identification, probable species identification	(++)	green
1.700 ... 1.999	probable genus identification	(+)	yellow
0.000 ... 1.699	not reliable identification	(-)	red

FIG. 1 Meaning of Bruker Biotyper score values.

No match avec d'autres espèces du groupe *scutellaris*!

Resistance (*Ae. aegypti*)

TABLEAU 2. NIVEAUX DE RESISTANCE DES LARVES DES NEUF POPULATIONS D'*AE. AEGYPTI* DE GUADELOUPE ET DE SAINT-MARTIN VIS-A-VIS DU TEMEPHOS ET DU MALATHION, PAR RAPPORT A LA SOUCHE SENSIBLE BORA-BORA.

Population	<i>Temephos</i>					
	DL50 (mg/l)	DL50 95%CI	RR50	DL95 (mg/l)	DL95 95%CI	RR95
Bora-Bora	0,008	0,008-0,009	1,00	0,014	0,012-0,017	1,00
LT	0,092	0,079-0,105	11,50	0,221	0,196-0,254	15,79
ABY	0,265	0,237-0,302	33,1	0,529	0,461-0,632	37,8
SF	0,123	0,111-0,136	15,4	0,233	0,211-0,262	16,6
BM	0,233	0,211-0,259	29,1	0,450	0,402-0,519	32,1
SXM (West)	0,071	0,062-0,081	8,9	0,160	0,143-0,183	11,4
SXM (East)	0,128	0,112-0,144	16,0	0,303	0,273-0,345	21,6
AB	0,132	0,115-0,152	16,5	0,557	0,440-0,763	39,8
DH	0,151	0,132-0,172	18,9	0,555	0,442-0,758	39,6
Bora-Bora	0,004	-	1,00	0,006	0,006-0,007	1,00
SBH	0,040	0,034-0,046	10,00	0,079	0,071-0,090	13,17




TABLEAU 3. STATUT DE RESISTANCE A LA DELTAMETHRINE DES FEMELLES ADULTES DES HUIT POPULATIONS D'*AE. AEGYPTI* DE GUADELOUPE, DE SAINT-MARTIN ET DE SAINT-BARTHELEMY.

Concentration Deltaméthrine - Temps d'exposition	Paramètres évalués	SBH	BT/GB	DH	AB	SF	SXM	ABY	BM
0,06% - 1h	Taux de KD	1	0,86	0,93	0,94	0,98	1	0,96	0,99
	Taux de mortalité après 24h	0,83	0,83	0,83	0,64	0,79	0,88	0,79	0,9
0,08% - 1h	Taux de KD	1	1	1	0,96	1	1	0,99	1
	Taux de mortalité après 24h	0,9	1	0,94	0,93	0,81	0,98	0,85	0,99
0,05% - 2h	KRR50 (IC 95%)	9,3 (9,1-9,5)	ND	17,7 (16,7-20,5)	28,1 (24,2-34,9)	13,7 (13,0-14,4)	12,4 (12,0-12,8)	8,0 (7,8-8,3)	13,6 (13,1-14,2)
0,05% - 2h	KRR99 (ci 95%)	7,5 (7,1-8,1)	ND	10,3 (8,4-19,3)	57,5 (37,9-105,6)	23,8 (19,8-29,8)	11,1 (9,8-13,1)	10,1 (9,2-11,1)	15,3 (13,4-17,9)

Resistance mutations: *Ae. aegypti* à Saint barth




FIGURE 2. FREQUENCE GENOTYPIQUE DES MUTATIONS KDR V1016I ET F1534C CHEZ LA SOUCHE DE REFERENCE BORA BORA ET HUIT POPULATIONS D'*AE. AEGYPTI* DE GUADELOUPE, DE SAINT MARTIN ET DE SAINT-BARTHELEMY.

Merci d'avoir suivi cette première journée !

Journées Scientifiques Annuelles d'Arbo-France

Jour 2 : Début 9H sur le 2nd lien

23-24 juin 2025, Val de Grâce - Paris